

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
30 January 2003 (30.01.2003)

PCT

(10) International Publication Number
WO 03/008591 A1

(51) International Patent Classification⁷: C12N 15/54

Jinwoo APT., #265, Samga-dong, 449-060 Yongin-city,
Kyonggi-do (KR).

(21) International Application Number: PCT/KR02/01344

(22) International Filing Date: 16 July 2002 (16.07.2002)

(74) Agent: LEE, Duck-Rog; 2nd Fl., Yeil Bldg., #700-19
Yorksam-dong, Kangnam-ku, 135-918 Seoul (KR).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
2001/42931 16 July 2001 (16.07.2001) KR

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): BIO HOLDINGS CO., LTD. [KR/KR]; 201-ho, Bioventure Center, c/o Korean Research Institute of Bioscience and Biotechnology, #52, Oun-dong, Yusong-gu, 305-333 Taejeon-city (KR).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant and

(72) Inventor: SUH, Joo-Won [KR/KR]; 8-ho, Myung-gimaeul, San 33-1, Nam-dong, 449-728 Yongin-city, Kyonggi-do (KR).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): YANG, Young-Yell [KR/KR]; 224-502, Hyojagreen APT., Jigok-dong, Nam-gu, 790-752 Pohang-city, Kyongsangbuk-do (KR). LEE, In-Hyung [KR/KR]; 32-906, Hansin 3 cha APT., #1-1, Banpo 2-dong, Seocho-gu, 137-042 Seoul (KR). KIM, Dong-Jin [KR/KR]; #97-1, Kyopyong-ri, Cheongsan-myun, 373-871 Ockcheon-gun, Chungcheongbuk-do (KR). HYUN, Chang-Gu [KR/KR]; 101-604,

Published:

- with international search report.
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ADENOSYLMETHIONINE SYNTHETASE FROM STREPTOMYCES SP., GENE SEQUENCES CODING THE SAME AND METHOD FOR MASS PRODUCTION OF SECONDARY METABOLITES INCLUDING ANTIBIOTICS THEREOF



(57) Abstract: Disclosed is an isolated nucleotide sequence encoding an enzyme catalyzing biosynthesis of SAM (SAM-s) and its amino acid sequence. Also, the present invention provides a method for mass production of a useful secondary metabolite including antibiotics using the isolated nucleotide sequence and SAM, where SAM acts as a methyl group donor.

WO 03/008591 A1

ADENOSYLMETHIONINE SYNTHETASE FROM STREPTOMYCES SP., GENE SEQUENCES CODING THE
SAME AND METHOD FOR MASS PRODUCTION OF SECONDARY METABOLITES INCLUDING ANTIBIOTICS
THEREOF

TECHNICAL FIELD

The present invention relates to S-adenosyl-L-methionine synthetase and a nucleotide sequence encoding the same, and more particularly, to an isolated nucleotide sequence encoding an enzyme catalyzing biosynthesis of SAM (SAM-s) from adenosyl triphosphate (ATP) and methionine, and its amino acid sequence. Also, the present invention is concerned with a method for mass production of a useful secondary metabolite including antibiotics using the nucleotide sequence and SAM.

10 PRIOR ART

S-adenosyl-L-methionine (SAM) is well known to play a critical role in cell growth and differentiation, essential for survival of living organisms including human beings. In living cells, SAM acts as a methyl group donor as well as a precursor for an aminopropyl group in a biosynthesis pathway of polyamine, where the methyl group and the polyamine are utilized in primary and secondary metabolisms.

15 It has been reported that SAM positively or negatively affects growth of bacteria including *E. coli* and *Bacillus subtilis*, thus causing their

life cycles to change in a manner of inhibiting cell growth or stimulating morphological differentiation.

In addition, the biological function of SAM is also found to be essential for primary and secondary metabolisms in plants and animals. Especially, it has been reported that SAM as a methyl group donor affects differentiation, causing morphological changes in plant or animal cells.

On the other hand, spectinomycin, which is an antibiotic derived from *Streptomyces spectabilis*, belongs to an aminoglycoside family and is composed of one sugar and two methyl groups originated from a methyl group donor, SAM.

DISCLOSURE OF THE INVENTION

Based on the fact that methyl groups of spectinomycin are derived from SAM, inventors of the present invention conducted intensive and thorough research into effects of SAM on biosynthesis of spectinomycin, resulting in the finding that SAM positively affects the biosynthesis of antibiotics, thereby increasing their production yield.

Therefore, it is an object of the present invention to provide an isolated nucleotide sequence encoding an enzyme catalyzing biosynthesis of SAM from *Streptomyces spectabilis* ATCC 27741 and an amino acid sequence translated from the isolated nucleotide sequence.

It is another object of the present invention to provide a method of increasing production of a useful secondary metabolite including antibiotics using SAM.

In accordance with the present invention, the first object is achieved by isolating a gene encoding an enzyme catalyzing SAM biosynthesis,

which is derived from *S. spectabilis*, by obtaining a PCR product of 4.0 kb from a gene library of *S. spectabilis* using PCR, and confirming presence of a gene of about 1.2 kb in the PCR product encoding an enzyme catalyzing SAM biosynthesis, by sequencing the PCR product assaying activity of its translational product.

In accordance with the present invention, the second object is achieved by producing SAM, which is synthesized by the translational product of the isolated nucleotide sequence or that which is commercially available, having an ability to stimulate production of an antibiotic.

BRIEF DESCRIPTION OF THE DRAWINGS

The above and other objects, features and other advantages of the present invention will be more clearly understood from the following detailed description taken in conjunction with the accompanying drawings, in which:

Fig. 1 is a restriction map of a gene encoding an enzyme catalyzing SAM biosynthesis (SAM-s) carried in a recombinant plasmid, pJWK0012, which is originated from an *E. coli* plasmid;

Fig. 2 is a result of a homology search comparing the amino acid sequence of "SAM-s" of the present invention to SAM synthetases from other microorganisms, obtained from GeneBank database, USA;

Fig. 3 is a graph showing an ability of "SAM-s" to synthesize SAM, using HPLC, where commercially available SAM is used as a control;

Fig. 4 is a graph showing an effect of SAM produced by "SAM-s" on production yield of actinorhodin in *S. lividans* TK23 transformed with

a gene encoding "SAM-s";

Fig. 5 is photograph showing an effect of SAM produced by "SAM-s" on production yield of actinorhodin in *S. lividans* TK23 treated with commercially available SAM; and

5 Fig. 6 is a graph showing an effect of SAM on production yield of undecylprodigiosin in *S. lividans* TK23.

BEST MODES FOR CARRYING OUT THE INVENTION

In accordance with the present invention, there is provided an enzyme catalyzing biosynthesis of S-adenosyl-L-methionine having an amino acid sequence consisting of the sequence shown in SEQ ID NO. 2,
10 which is derived from *Streptomyces spectabilis* ATCC 27741.

In accordance with the present invention, there is provided an isolated nucleotide sequence consisting of the sequence shown in SEQ ID NO. 1, which encodes the amino acid sequence of SEQ ID NO. 2.

15 In accordance with the present invention, there is also provided a method of producing a useful secondary metabolite including antibiotics in a *Streptomyces* species-originated transformant into which a gene encoding an enzyme catalyzing biosynthesis of SAM is introduced, thereby improving productivity of the useful secondary metabolite.

20 In accordance with the present invention, there is further provided a method of producing a useful secondary metabolite including antibiotics by directly adding SAM to culture medium containing antibiotic-producing bacteria, thereby improving productivity of the useful secondary metabolite.

In the present invention, preferable examples of the useful secondary metabolite include antibiotics, such as polyketide antibiotics, anti-cancer agents, and vermicides.

5 In the present invention, a gene encoding an enzyme catalyzing SAM biosynthesis (SAM-s) is isolated from cosmid clones containing genes encoding enzymes stimulating biosynthesis of spectinomycin, which is obtained from a cDNA library of *S. spectabilis* ATCC 27741, where a 3.9 kb clone is primarily obtained. The nucleotide sequence of the 3.9 kb clone is partially determined by performing nucleotide
10 sequencing, and its homology to known SAM synthetases, which mediate biosynthesis of SAM using ATP and methionine as substrates, is examined, based on the obtained nucleotide sequence, indicating the possible presence of a gene encoding an enzyme catalyzing SAM biosynthesis. The 3.9 kb clone is demonstrated to carry the gene
15 encoding the enzyme catalyzing SAM biosynthesis by in vitro assaying the activity of its translated product.

In accordance with the present invention, a portion of the 3.9 kb clone, containing the gene encoding an enzyme catalyzing SAM biosynthesis (SAM-s), is introduced into *Streptomyces* species to assay
20 production yield of an antibiotic, actinorhodin, and also, the effect of SAM on production of antibiotics is further investigated through direct treatment of cells with SAM, thereby demonstrating that SAM is effective in improving productivity of secondary metabolites including antibiotics.

In embodiments of the present invention, *Streptomyces lividans*
25 TK23, which is commercially available, is transformed with the gene encoding "SAM-s", and the resulting transformant exhibits mass production of actinorhodin, demonstrating that SAM is effective in

enhancing antibiotic productivity of cells. Herein, application of SAM for high production of antibiotics is not limited to the transformant and the antibiotic, actinorhodin, but the effectiveness of SAM on production of antibiotics can be achieved with all *Streptomyces* species transformed with
5 a gene encoding an enzyme catalyzing SAM biosynthesis.

In addition, when SAM is directly added to culture medium containing antibiotic-producing bacteria, productivity of antibiotics is increased 5 to 10 times, and especially, production of polyketide antibiotics is significantly increased.

10 The present invention will be explained in more detail with reference to the following examples in conjunction with the accompanying drawings. However, the following examples are provided only to illustrate the present invention, and the present invention is not limited to them.

15 **EXAMPLE 1 : Cloning of a gene encoding an enzyme capable of SAM biosynthesis from cosmid clones of *Streptomyces spectabilis* ATCC 27741**

Genes encoding enzymes catalyzing biosynthesis of an antibiotic are typically located together in a specific region of a genome. Therefore, there was used two cosmid clones harboring 30 to 40 kb
20 fragment, which carries a gene family consisting of genes encoding enzymes participating in spectinomycin biosynthesis and may also include a gene encoding methyltransferase enzyme, one of enzymes mediating spectinomycin biosynthesis, which functions to transfer methyl groups. After digestion of the two cosmid clones with restriction enzymes,

Southern Blotting was performed using metK gene, having high homology to methyltransferase at the nucleotide sequence level, as a probe.

As a result of Southern Blotting, a positive spot was observed, indicating a 3.9 kb fragment inserted into a BamHI site of pHCG121.

5 3.9 kb fragment was then subcloned into a BamHI site of pBluescript KS(+), giving a recombinant plasmid pHCG1647. From the subcloned 3.9 kb fragment, a 2.5 kb fragment, which is believed to carry a gene encoding an enzyme catalyzing SAM biosynthesis, was subcloned again into pBluescript KS(+) to form a recombinant plasmid pJWK0012.

10 **EXAMPLE 2 : Determination of nucleotide sequence of the cloned 2.5 kb fragment and its corresponding amino acid sequence**

In order to determine a nucleotide sequence of the cloned 2.5 kb fragment and its corresponding amino acid sequence, the 2.5 kb insert carried in pJWK0012 prepared in the Example 1 was digested with
15 restriction enzymes, ApaI, SalI and SacI, and then subcloned, followed by nucleotide sequencing. Fig. 1 shows a restriction map of the 2.5 kb fragment in pJWK0012 and its translational orientation.

Based on the nucleotide sequence of the 2.5 kb fragment, its amino acid sequence was obtained through search using a Codon Preference
20 program (Bibb, M. J. et al., Gene, 1984). As a result, the 2.5 kb fragment was found to have an open reading frame consisting of a coding region ranging from nt 835 to nt 2051, which may express a protein consisting of 464 amino acid. The translational product of the open reading frame was, in the present invention, called "SAM-s".

To investigate the homology of "SAM-s" to other known proteins, the amino acid sequence of "SAM-s" was compared to those of SAM synthetases of *Streptomyces coelicolor*, *Bacillus subtilis* and *Escherichia coli*, which were obtained from GeneBank DataBase (USA). With reference to Fig. 2, it was found that "SAM-s" shares high homology with other synthetases. Also, "SAM-s" of the present invention was found to have homology to some methyltransferases from microorganisms.

EXAMPLE 3 : Assay for activity of "SAM-s"

In order to analyze activity of "SAM-s", the gene encoding "SAM-s" was expressed in *E. coli*, and the resulting translational product, "SAM-s", was then isolated.

To express the gene encoding "SAM-s" in *E. coli*, the gene was inserted into a pET-21a vector, and then introduced into *E. coli* BL21. The expressed gene product, "SAM-s" was isolated using a His-Tag purification system. Thereafter, 10 to 50 μ l of the enzyme solution containing the protein "SAM-s" was added to a reaction mixture containing 100 mM of Tris-HCl, 200 mM of KCl, 10 mM of $MgCl_2$, 1 mM DTT, 5 mM ATP, and 5 mM methionine, followed by incubation for 120 min at 30 °C. After the incubation, reaction products were analyzed through HPLC using Reverse C18 column. In this regard, the column loaded with sample was initially equilibrated with a solution of 0.1 M of NaH_2PO_4 /acetonitrile at a ratio of 98:2 (V/V), pH 2.65. Then, a second solution comprising 0.15 M NaH_2PO_4 /acetonitrile at a ratio of 74:26 (V/V) was applied with continuous mixing with the first solution, forming a concentration gradient.

As shown in Fig. 3, the product of the catalytic activity of the protein expressed in *E. coli* is proven to be SAM. That is, when the expressed protein is supplied with ATP and methionine as substrates, the product has an HPLC retention time identical to commercially available SAM, indicating that the protein expressed in *E. coli* has an activity to synthesize SAM using ATP and methionine as substrates.

EXAMPLE 4 : Effect of in vivo-synthesized SAM on productivity of actinorhodin in *S. lividans* TK23

The gene encoding "SAM-s" was first inserted into pWHM3, which is a shuttle vector between *E. coli* and *Streptomyces* species, giving an expression vector pSAM-s. The plasmid pSAM-s was then introduced into *S. lividans* TK23. The resulting transformant, *Streptomyces lividans* TK-23 harboring pSAM-s, was deposited in the Korean Culture Center of Microorganisms with accession No. KCCM 10397 on July 2, 2002. The transformant was incubated in one liter of a medium including 50 g of glycerol, 5 g of glutamic acid, 21 g of morpholinopropane sulfonic acid, 200 mg of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 100 mg of $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 100 mg NaCl, 82 mg KH_2PO_4 , 9 mg $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, and 2 ml of trace element solution, adjusted to pH 6.5. During incubation for 7 days at 28 °C, production yield of actinorhodin, which is a main antibiotic produced from *S. lividans* TK23, was analyzed. The results are shown in Fig. 4.

As apparent in Fig. 4, when SAM was over-produced in *S. lividans* TK23 through over-expression of "SAM-s", it was observed that production of actinorhodin in the transformant was enhanced to over six times in comparison with that of a wild type *S. lividans*.

EXAMPLE 5 : Effect of externally added SAM on productivity of actinorhodin

Based on the finding that in vivo over-expressed SAM positively affects production yield of actinorhodin in *S. lividans* TK23, an effect of SAM on productivity of actinorhodin was investigated when commercially available SAM is added directly to culture medium containing *S. lividans* TK23. As such, wild type *S. lividans* TK23 was treated with 1 mM of commercially available SAM.

The result is shown in Fig. 5, where actinorhodin produced in *S. lividans* TK23 treated with SAM, and the control not treated with SAM, indicated by a blue color. As shown in Fig. 5, it was found that *S. lividans* TK23 treated with SAM produces more actinorhodin than *S. lividans* TK23 not treated with SAM, demonstrating that SAM positively affects productivity of actinorhodin.

EXAMPLE 6 : Effect of SAM on productivity of undecylprodigiosin in *S. lividans* TK23

S. lividans TK23 transformed with the vector pSAM-s was incubated under the same culture conditions as those used for production of actinorhodin. To determine the amount of undecylprodigiosin produced, after adjusting pH to 12, absorbance was measured at 468 nm, and concentration of the antibiotic was calculated according to the following formula: concentration = OD value \times 9.4673.

As apparent Fig. 6, productivity of undecylprodigiosin was very high in comparison with a control not transformed with the vector pSAM-

s, indicating that SAM positively affects production yield of undecylprodigiosin.

EXAMPLES 7 to 13: Effect of SAM on productivity of antibiotics in *Streptomyces* species

5

TABLE 1
Culture medium and culture condition

Exp.	Antibiotic	<i>Streptomyces</i> sp.	Culture medium (/L) and culture condition
7	Avermectin	<i>S. avermitilis</i>	15g glucose, 0.5g asparagine, 0.5g K ₂ HPO ₄ , pH 7.0, 25 °C, incubation for 5 days
8	Monensin	<i>S. cinnamonensis</i>	2.5% glucose, 1.5% soybean meal, 0.3% CaCO ₃ , 0.03% FeSO ₄ 7H ₂ O, 0.003% MnCl ₂ 4H ₂ O, pH 7.0, 30 °C, incubation for 5 days
9	Spectinomycin	<i>S. spectabilis</i>	10g Maltose, 5g tryptone, 1g K ₂ HPO ₄ , 2g NaCl, pH7.0, 30 °C, incubation for 5 days
10	Doxorubicin	<i>S. peucetius</i>	60g Glucose, 8g yeast extract, 20g malt extract, 2g NaCl, 15g MOPS sodium salt, 0.1g MgSO ₄ , 0.01g FeSO ₄ 7H ₂ O, 0.01g ZnSO ₄ 7H ₂ O, pH 7.0, 30 °C, incubation for 5 days
11	Streptomycin	<i>S. griseus</i>	1 % glucose, 0.4 % peptone, 0.2 % meat extract, 0.2 % yeast extract, 0.5 % NaCl, 0.025 % MgSO ₄ 7H ₂ O, pH 7, 30 °C, incubation for 5 days
12	Tetracyclin	<i>S. aureofaciens</i>	3 % corn flour, 4 % corn steep liquor, 5 % corn starch, 0.7 % (NH ₄) ₂ SO ₄ , 0.1 % NH ₄ Cl, 5 ppm CoCl ₂ , 0.9 % CoSO ₃ , 2 % rice bran oil, pH7, 28 °C, incubation

			for 5 days
13	Chlortetracyclin	<i>S. aureofaciens</i>	1 % sucrose, 1 % corn steep liquor, 0.2 % $(\text{NH}_4)_2\text{HPO}_4$, 0.2 % KH_2PO_4 , 0.1 % CaCO_3 , 0.025 % $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.005 % $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 0.00033 % $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.00033 % $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$, incubation for 5 days

Each *Streptomyces* species was incubated in its corresponding culture medium according to Table 1, and treated with 1 mM of SAM (Sigma, USA). After incubation for 5 days, antibiotic concentration was measured in each culture medium. The results are given in Table 2, below. It was found that each *Streptomyces* species, as a treatment group, produces a much higher amount of its specific antibiotic than a control group not treated with SAM.

TABLE 2

Production amount of an antibiotic in *Streptomyces* sp. treated or not treated with SAM

Exp.	Antibiotic	<i>Streptomyces</i> sp.	Production amount of a control group ($\mu\text{g/ml}$)	Production amount of a treatment group ($\mu\text{g/ml}$)
7	Avermectin	<i>S. avermitilis</i>	5	25
8	Monensin	<i>S. cinnamomensis</i>	30	180
9	Spectinomycin	<i>S. spectabilis</i>	5	35
10	Doxorubicin	<i>S. peucetius</i>	38	300
11	Streptomycin	<i>S. griseus</i>	101	602
12	Tetracycline	<i>S. aureofaciens</i>	30	188

13

13	Chlortetracycline	<i>S.</i> <i>aureofaciens</i>	25	130
----	-------------------	----------------------------------	----	-----

As shown in Table 2, *Streptomyces* species treated with SAM produced 5 to 10 times more antibiotic than the control, indicating that SAM positively affects production yield of various antibiotics.

INDUSTRIAL APPLICABILITY

5 As described hereinbefore, the present invention provides an isolated nucleotide sequence of a gene encoding an enzyme catalyzing biosynthesis of SAM, which is derived from *Streptomyces spectabilis* ATCC 27741, and its amino acid sequence. SAM, which is produced by the enzyme of the present invention or purchased commercially, is
10 very effective in increasing productivity of various antibiotics. Therefore, the isolated nucleotide sequence of the present invention is capable of being utilized in mass production of secondary metabolites including antibiotics, and thus is very useful in pharmaceutical industries.


15

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

To: Suh, Joo-Won
San 38-2 Namdong, Yongin,
Kyunggi-do, 449-728, Korea

RECEIPT IN THE CASE OF AN ORIGINAL
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR : <i>Streptomyces lividans</i> TK-23 harboring pSAM-s	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM-10397
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: <input type="checkbox"/> a scientific description <input type="checkbox"/> a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on Jul. 2, 2002. (date of the original deposit) ¹	
IV. INTERNATIONAL DEPOSITARY AUTHORITY	
Name : Korean Culture Center of Microorganisms Address : 361-221, Yurim R/D Hongje-1-dong, Seodaemun-gu SEOUL 120-091 Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s): Date: Jul. 9, 2002 <div style="text-align: center; margin-top: 10px;">  </div>

¹ Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired: where a deposit made outside the Budapest Treaty after the acquisition of the status of international depositary authority is converted into a deposit under the Budapest Treaty, such date is the date on which the microorganism was received by the international depositary authority.

CLAIMS

1. S-adenosyl-L-methionine synthetase, having an amino acid sequence consisting of the sequence shown in SEQ ID NO. 2, which is derived from *Streptomyces spectabilis* ATCC 27741.

5 2. A nucleotide sequence of SEQ ID NO. 1, encoding the amino acid sequence of claim 1.

3. A method of producing a useful secondary metabolite including antibiotics using a transformant into which a gene encoding S-adenosyl-L-methionine synthetase is introduced, thereby improving productivity of
10 said useful secondary metabolite.

4. A method of producing a useful secondary metabolite including antibiotics comprising adding S-adenosyl-L-methionine to culture medium containing antibiotic-producing bacteria, thereby improving productivity of said useful secondary metabolite.

1/4

FIG. 1





FIG. 2

	1	50
Mycobacterium	MSEKGRLFIS	ESVTEGHPDK ICDAISDSVL DALLAADPRS RVAVETLVTI GQVEVVG EYT ISAKEAFADI INTVRARILE IGYDSSDKGF DGAICGVNIG
Streptomyces	..MSRRLFIS	ESVTEGHPDK IADQISDTIL DALLREDPTS RVAVETLITIT GLVHVAGEYT T...KAYAPI AQLVREKILE IGYDSSKKGF DGASCGSVS
Escherichia	..MAKELFIS	ESVSEGHDPK IADQISDAVL DAILEQDPKA RVACETYVKI GNLVSGEIT T...SAVVDI EEITRNTVRE IGYVESDMGF DANSCAVLSA
Bacillus	MSKNRRLFIS	ESVTEGHPDK ICDQISDSIL DEILKEDPNA RVACETSVIT GLVLSGEIT T...STYVDI PKTVRQTIKE IGYTRAKYGF DAETCAVLTS
Staphylococcus	MLNKRRLFIS	ESVTEGHPDK IADQVSDAIL DAILKDDPNA RVACETITV GHALIAGEIS T...ITYVDI PKVRETIKE IGYTRAKYGY DYETHAILTI
Consensusrlfts	esvtEGHPDK.lcDq!SD.!L DaLL..DP.s RVA.ET.!tT Glvll!GE!t t....a..di ...vR...!E IGY..sd.GX da.sca!..a
	101	150
Mycobacterium	IGAQSPDIAQ	GVDTAHEARY EG.AADPLDS QGAGDQGLMF GYAINAPEL MPLPIALAER LSRLTEVRK NGVLPYLRLPD GKQVTIAYE DN.VPVRLDI
Streptomyces	IGAQSPDIAQ	GVDTAYESRV EG.DEELDR QGAGDQGLMF GYACDEIPEL MPLPIELAER LSRLSEVRK NGTIPYLRLPD GKQVTIEYD GD.KAVRLDI
Escherichia	ICKQSPDINQ	GVDRADPLEQ GAGDQGLMF GYATNETDVL NPAPITYAER LVQQAQEVK NGTLPVLRPD AKSQVTIFYD DG.KIVGIDA
Bacillus	IDEQSADIAM	GVDQALEARE GINSDEEIEA IGAGDQGLMF GYACNETKEL MPLPISLARK LARLSEVRK EDILPYLRPD GKQVTIVEYD ENNKPVRLDA
Staphylococcus	IDEQSPDIAQ	GVDKALEYRD KD.SEEIEIA TGAGDQGLMF GYATNETETY MPLAIYLSHQ LAKRLSDVRK DGTILNYLRPD GKQVTIVEYD ENDNPVRID
Consensus	Ig.QSpDlaq	Gvd.A.a.r.deld. GAGDQGLMF GYAc#sTpel MPLpI.laEr l...Rls#VRK ns.lpyLRPD gKtQvtI.Y# d...pVrIdt
	201	250
Mycobacterium	VVISTQHA..	..DIDLEKTL DPDIREFVYN TVLDDLAHEI LDAST..VRVLNPIGKFLV GGPNGDAGLT GRKIIIDTYG GVARHGGGAF SGKDPKVDK
Streptomyces	VVSSQHAS..	..DIDLESLL APDIREFVVE PELKALVEDG IKLETEGYRLLVNPITGRFEI GGPNGDAGLT GRKIIIDTYG GMSREGGGAF SGKDPKVDK
Escherichia	VVLSTQSE..	..EIDQKS.L QEAVMEEIIR PILP...AEV LTSAT...KFFINPTGRFVI GGPNGDCGLT GRKIIIDTYG GVARHGGGAF SGKDPKVDK
Bacillus	IVISTQHP..	..EITLQ.I QRNIKEHVIN PWP...EEL IDEET...KYFINPTGRFVI GGPQGDAGLT GRKIIIDTYG GVARHGGGAF SGKDPKVDK
Staphylococcus	IVVSTQHA..	..DVTLEQ.I QEDIKAEVIY PTVP...ENL INEQT...K YINPTGRFVI GGPQGDAGLT GRKIIIDTYG GIARHGGGCF SGKDPKVDK
Consensus	IviStQh.e.	..#!dle..l q...!e.v... pvl.....#.T...rf.!NPTGrFvl GGPmGdGLT GRKIIIDTYG G.aREGGGaF SGKDPsKVDR
	301	350
Mycobacterium	SAAYANRVVA	KNVVAAGLAE RVEQVAYAI GKAAPVGLFY ETFGTETEDPVKIEKAIGEY FDLRPGAIIR DLNLLRPIYAPTA AYGHFGRITD
Streptomyces	SAAYANRVVA	KNVVAAGLAS RCEVQVAYAI GKAEPVGLFY ETFGTNTIDIDKIEQAISEY FDLRPAAIIR SLDLLRPIYSQTA AYGHFGRSLP
Escherichia	SAAYAARYVA	KNIVAAGLAD RCEIQVSYAI GVAEPTSINV ETFGTEKVPSEQLTLLVREF FDLRPYGLIQ MLDLLEPIYKETA AYGHFGREH.
Bacillus	SAAYAARYVA	KNIVAAGLAD RCEVQLAYAI GVAQPVSIISI NTFGSGKASEEKLIEVVRNN FDLRPAGIIR MLDLRRPIYKOTA AYGHFGRBDV
Staphylococcus	SAAYAARYVA	KNIVAAGLAD RCEVQLAYAI GVAEPVSIISI DTFGTCKVS QQLVEAVRKE FDLRPAGIIR MLDLRRPIYKOTA AYGHFGRITD
Synechocystis	SAAYAARYVA	KNIVAAGLAD RCEVQVSYAI GVARPVSVLI DTFGTCKVDEekl...!e. F#LRPagiI. .L#LI...piY.ta AYGHFGR.d.
Consensus	SAAYaArYVA	KN!VAaGLad rcE!GvaYAI GvA.Pvs..! #TFGt.kvd.


2/4

FIG. 3



3/4

FIG. 4



[Note]

- A : Cell growth
- B : Production of antibiotics
- : Transformant
- ◆ : Wild type.

4/4

FIG. 5





FIG. 6



[Note] ■ : Treatment group
◆ : Control group

SEQUENCE LISTING

<110> SUH, Joo-Won

<120> Adenosylmethionine synthetase from Streptomyces sp. and gene sequences coding the same

<130> YL02007PCT

<140> PCT/KR02/01344

<141> 2002-07-16

<150> KR 2001-42931

<151> 2001-07-16

<160> 2

<170> PatentIn version 3.1

<210> 1

<211> 2349

<212> DNA

<213> Streptomyces spectabilis ATCC 27741

<220>

<221> CDS

<222> (837)..(2096)

<223>

<400> 1

```

gggcccgcgt cagctcgtc gcgcggaaca ccgctgcc ggacccggcg ggcgtcgacc      60
tcgtgcccggt gggcaccgcc gtacagctcc gcgaggccgt cctgaaggcg gccgccgacg      120
ccgacgcggt ggtgatggcc gccccgtgg ccgacttccg cccggcccgcg tacgcccccg      180
gaaagatcaa gaagcaggac ggcgaggacc cggccccgct caccctcgtc cgcaatccgg      240
atacctcgc cgagctctcg ggcgaccgcc cgcgccccgg ccaggtcgtc gtcggcttcg      300
cggccgagac ggacaacgtg ctggccaacg gcgggccaag ctgcccgcga aggctgcgac      360
ctcctcgtag tgaacgaagt cggtgagcgc aagatgttcg gctctgagcg gaacgaggcc      420
gtggtgctcg gtgccgacgg cagtgagacg ccggtgccct acggcccca ggacgatctc      480
gccgacacgg tctgggatct ggtcgaccg cgccttgacc ggtcgtaaca gcctgaggc      540
catcggtcg gatcgtggcc gggccgatgg ccggaaccgg ccccggcgcg ccatgcgcc      600
ctccaggctc tctaaacca ccctaaacc gccaaatcg ggccttgagc cctctgttg      660
agaccgatcg tcgtatcgca caatgcagt cgcaggtca caggcctctc ttctgacgag      720
agaggtggcc atcgggccgt gtgtgaccga taaactggc tcggacgtcg tcgagcgag      780
ctctcgcccc gtccatcaat gatcagccag cagccgtgc aaccacaggg agcgtg atg      839
Met

```

tcc cgc cgt ctc ttc acc tcg gag tcc gtg acc gag ggt cac ccc gac Ser Arg Arg Leu Phe Thr Ser Glu Ser Val Thr Glu Gly His Pro Asp 5 10 15	887
aag atc gct gac cag atc agc gac acc att ctc gac gcg ctt ctg cgt Lys Ile Ala Asp Gln Ile Ser Asp Thr Ile Leu Asp Ala Leu Leu Arg 20 25 30	935
gag gac ccg acg tcc cgg gtc gcc gtc gaa acg ctc atc acc acc ggc Glu Asp Pro Thr Ser Arg Val Ala Val Glu Thr Leu Ile Thr Thr Gly 35 40 45	983
ctc gtg cac gtc gcc ggt gag gtc acg acc aag gcg tac gcg ccg atc Leu Val His Val Ala Gly Glu Val Thr Thr Lys Ala Tyr Ala Pro Ile 50 55 60 65	1031
gcg cag ctg gtg cgc gag aag atc ctc gag atc gga tac gac tcc tcg Ala Gln Leu Val Arg Glu Lys Ile Leu Glu Ile Gly Tyr Asp Ser Ser 70 75 80	1079
aag aag ggc ttc gac ggc gcc tcc tgc ggc gtc tcg gtg tcc atc ggc Lys Lys Gly Phe Asp Gly Ala Ser Cys Gly Val Ser Val Ser Ile Gly 85 90 95	1127
gcg cag tcc ccg gac atc gcg cag ggc gtc gac acg gcg tac gag agc Ala Gln Ser Pro Asp Ile Ala Gln Gly Val Asp Thr Ala Tyr Glu Ser 100 105 110	1175
cgt gtc gag ggc gac gag gac gag ctc gac cgg cag ggc gcc ggt gac Arg Val Glu Gly Asp Glu Asp Glu Leu Asp Arg Gln Gly Ala Gly Asp 115 120 125	1223
cag ggc ctg atg ttc ggc tac gcc tgc gac gag acc ccg gag ctg atg Gln Gly Leu Met Phe Gly Tyr Ala Cys Asp Glu Thr Pro Glu Leu Met 130 135 140 145	1271
ccg ctc ccg atc cac ctc gcg cac cgc ctc tcg cgc cgc ctc tcc gag Pro Leu Pro Ile His Leu Ala His Arg Leu Ser Arg Arg Leu Ser Glu 150 155 160	1319
gtg cgc aag aac ggg acc atc ccc tac ctg cgc ccc gac ggc aag acc Val Arg Lys Asn Gly Thr Ile Pro Tyr Leu Arg Pro Asp Gly Lys Thr 165 170 175	1367
cag gtc acc atc gag tac gac ggc gac aag gcc gtc cgc ctc gac acg Gln Val Thr Ile Glu Tyr Asp Gly Asp Lys Ala Val Arg Leu Asp Thr 180 185 190	1415
gtc gtc gtc tcc tcg cag cac gcg tcg gac atc gac ctg gag tcg ctg Val Val Val Ser Ser Gln His Ala Ser Asp Ile Asp Leu Glu Ser Leu 195 200 205	1463
ctc gcc ccc gac atc cgc gag ttc gtc gtc gag ccg gag ctc aag gcc Leu Ala Pro Asp Ile Arg Glu Phe Val Val Glu Pro Glu Leu Lys Ala 210 215 220 225	1511

ctg gtc gag gac ggc atc aag ctg gtc gtc gag ccg gag ctc aag gcc Leu Val Glu Asp Gly Ile Lys Leu Val Val Glu Pro Glu Leu Lys Ala 230 235 240	1559
ctg gtc gag gac ggc atc aag ctg gag acc gag ggc tac cgc ctc ctg Leu Val Glu Asp Gly Ile Lys Leu Glu Thr Glu Gly Tyr Arg Leu Leu 245 250 255	1607
gtc aac ccg acc ggc cgc ttc gag atc ggc ggc ccg atg ggt gac gcg Val Asn Pro Thr Gly Arg Phe Glu Ile Gly Gly Pro Met Gly Asp Ala 260 265 270	1655
ggc ctg acc ggc cgc aag atc atc atc gac acc tac ggc ggc atg tcc Gly Leu Thr Gly Arg Lys Ile Ile Ile Asp Thr Tyr Gly Gly Met Ser 275 280 285	1703
cgc cac ggc ggc ggc gcc ttc tcc ggc aag gac ccg tcc aag gtg gac Arg His Gly Gly Gly Ala Phe Ser Gly Lys Asp Pro Ser Lys Val Asp 290 295 300 305	1751
cgc tcg gcg gcg tac gcg atg cgc tgg gtc gcc aag aac gtg gtg gcc Arg Ser Ala Ala Tyr Ala Met Arg Trp Val Ala Lys Asn Val Val Ala 310 315 320	1799
gcg ggc ctc gcc tcg cgc tgc gag gtc cag gtc gcc tac gcc atc ggc Ala Gly Leu Ala Ser Arg Cys Glu Val Gln Val Ala Tyr Ala Ile Gly 325 330 335	1847
aag gcc gag ccg gtc ggt ctg ttc gtc gag acc ttc ggc acc aac acg Lys Ala Glu Pro Val Gly Leu Phe Val Glu Thr Phe Gly Thr Asn Thr 340 345 350	1895
atc gac acg gac aag atc gag cag gcc atc agc gag gtc ttc gac ctc Ile Asp Thr Asp Lys Ile Glu Gln Ala Ile Ser Glu Val Phe Asp Leu 355 360 365	1943
cgc ccg gcc gcg atc atc cgc agc ctc gac ctg ctc cgc ccg atc tac Arg Pro Ala Ala Ile Ile Arg Ser Leu Asp Leu Leu Arg Pro Ile Tyr 370 375 380 385	1991
tcc cag acc gcc gcg tac ggc cac ttc ggc cgc tcg ctg ccg gag ttc Ser Gln Thr Ala Ala Tyr Gly His Phe Gly Arg Ser Leu Pro Glu Phe 390 395 400	2039
acc tgg gag aag acg gac cgc gtg gac gcg ctg ccg aag gcc gcc ggt Thr Trp Glu Lys Thr Asp Arg Val Asp Ala Leu Arg Lys Ala Ala Gly 405 410 415	2087
ctg gag agc tgatctcctg ccgcttggtc actgaggccg tgccccctcaa Leu Glu Ser 420	2136
ggggcaccgg gcctcagtgc gtcagctgct ggagccgttc ggcatcgaca cgtcgtcgcc	2196
ctcgtcgggg tcttgaggcc cgaagcgccg ccggatgtcg tcaacgacat cgggcttgct	2256
gtgggcgtgg gcgagcttga tgagccgctc gacgaacagg tcgcgttccg gggcctcggg	2316

ggagatgccg aaccatttca ggaaggcgtc gac

2349

<210> 2

<211> 420

<212> PRT

<213> Steptomyces spectabilis ATCC 27741

<400> 2

Met Ser Arg Arg Leu Phe Thr Ser Glu Ser Val Thr Glu Gly His Pro
 1 5 10 15

Asp Lys Ile Ala Asp Gln Ile Ser Asp Thr Ile Leu Asp Ala Leu Leu
 20 25 30

Arg Glu Asp Pro Thr Ser Arg Val Ala Val Glu Thr Leu Ile Thr Thr
 35 40 45

Gly Leu Val His Val Ala Gly Glu Val Thr Thr Lys Ala Tyr Ala Pro
 50 55 60

Ile Ala Gln Leu Val Arg Glu Lys Ile Leu Glu Ile Gly Tyr Asp Ser
 65 70 75 80

Ser Lys Lys Gly Phe Asp Gly Ala Ser Cys Gly Val Ser Val Ser Ile
 85 90 95

Gly Ala Gln Ser Pro Asp Ile Ala Gln Gly Val Asp Thr Ala Tyr Glu
 100 105 110

Ser Arg Val Glu Gly Asp Glu Asp Glu Leu Asp Arg Gln Gly Ala Gly
 115 120 125

Asp Gln Gly Leu Met Phe Gly Tyr Ala Cys Asp Glu Thr Pro Glu Leu
 130 135 140

Met Pro Leu Pro Ile His Leu Ala His Arg Leu Ser Arg Arg Leu Ser
 145 150 155 160

Glu Val Arg Lys Asn Gly Thr Ile Pro Tyr Leu Arg Pro Asp Gly Lys
 165 170 175

Thr Gln Val Thr Ile Glu Tyr Asp Gly Asp Lys Ala Val Arg Leu Asp
 180 185 190

Thr Val Val Val Ser Ser Gln His Ala Ser Asp Ile Asp Leu Glu Ser
 195 200 205
 Leu Leu Ala Pro Asp Ile Arg Glu Phe Val Val Glu Pro Glu Leu Lys
 210 215 220
 Ala Leu Val Glu Asp Gly Ile Lys Leu Val Val Glu Pro Glu Leu Lys
 225 230 235 240
 Ala Leu Val Glu Asp Gly Ile Lys Leu Glu Thr Glu Gly Tyr Arg Leu
 245 250 255
 Leu Val Asn Pro Thr Gly Arg Phe Glu Ile Gly Gly Pro Met Gly Asp
 260 265 270
 Ala Gly Leu Thr Gly Arg Lys Ile Ile Ile Asp Thr Tyr Gly Gly Met
 275 280 285
 Ser Arg His Gly Gly Gly Ala Phe Ser Gly Lys Asp Pro Ser Lys Val
 290 295 300
 Asp Arg Ser Ala Ala Tyr Ala Met Arg Trp Val Ala Lys Asn Val Val
 305 310 315 320
 Ala Ala Gly Leu Ala Ser Arg Cys Glu Val Gln Val Ala Tyr Ala Ile
 325 330 335
 Gly Lys Ala Glu Pro Val Gly Leu Phe Val Glu Thr Phe Gly Thr Asn
 340 345 350
 Thr Ile Asp Thr Asp Lys Ile Glu Gln Ala Ile Ser Glu Val Phe Asp
 355 360 365
 Leu Arg Pro Ala Ala Ile Ile Arg Ser Leu Asp Leu Leu Arg Pro Ile
 370 375 380
 Tyr Ser Gln Thr Ala Ala Tyr Gly His Phe Gly Arg Ser Leu Pro Glu
 385 390 395 400
 Phe Thr Trp Glu Lys Thr Asp Arg Val Asp Ala Leu Arg Lys Ala Ala
 405 410 415
 Gly Leu Glu Ser
 420

INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR02/01344

A. CLASSIFICATION OF SUBJECT MATTER

IPC7 C12N 15/54

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7 C12N 15/54, C12N 9/10

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CA, PubMed, Blast, Delphion, "S-adenosyl-L-methionine synthetase", "Streptomyces", "S-adenosyl-L-methionine", "antibiotic"

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	NCBI Accession # AF117274, 31 Mar. 1999.	1, 2
X	JP 09-224690 A2 (Shiseido Co., Ltd. & Takeda Chem. Ind., Ltd.), 02 Sep. 1997. See abstract.	3, 4
Y	Seno, E.T. et al., Antimicrob. Agents Chemother., 21(5), 758-63, 1982. See abstract.	3, 4
Y	Merali, S. et al., Proc. Natl. Acad. Sci. USA, 96(5), 2402-7, 1999. See abstract.	4

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

- * Special categories of cited documents:
- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed
- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

25 NOVEMBER 2002 (25.11.2002)

Date of mailing of the international search report

25 NOVEMBER 2002 (25.11.2002)

Name and mailing address of the ISA/KR

Korean Intellectual Property Office
920 Dunsan-dong, Seo-gu, Daejeon 302-701,
Republic of Korea

Facsimile No. 82-42-472-7140

Authorized officer

LEE, Cheo Young

Telephone No. 82-42-481-5594



INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/KR02/01344

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
JP 09-224690 A2	02 Sep. 1997	none	